

***Rhodotorula toruloides* for carotenoid production using waste hardwood biomass**

Stefano Bertacchi,^{a,b} Francesca Sabatini,^{b,c} Giovanni Maria Bernardini^a, Matilde Dameri^a, Veronica Termopoli,^{b,c} Danilo Porro^{a,b}, Marco Orlandi,^{b,c} Heiko Lange,^{b,c,d,*} Paola Branduardi^{a,b,*}

E-mail: stefano.bertacchi@unimib.it

^a Department of Biotechnology and Biosciences, University of Milano-Bicocca, 20126 Milan, Italy

^b NBFC – National Biodiversity Future Center, 90133 Palermo, Italy

^c Department of Earth and Environmental Sciences, University of Milano-Bicocca, 20126 Milan, Italy

^d Biochemical Process Engineering, Division of Chemical Engineering, Department of Civil, Environmental and Natural Resources Engineering, Luleå University of Technology, SE-971 87 Luleå, Sweden

Keywords: *Rhodotorula toruloides*, Carotenoids, Biodiversity valorisation, Biorefinery, Lignocellulosic biomass, Enzymatic hydrolysis

Abstract:

This work explores the potential of underutilized urban pruning residues from hardwood as feedstocks for bioprocesses based on the carotenogenic yeast *Rhodotorula toruloides*, investigating the correlation between biomass composition and carotenoids production. Enzymatic hydrolysates from woods and barks of sessile oak and mulberry tree were used as substrates for microbial fermentation, obtaining superior titers and productivity of β -carotene and torulene when compared to previous published processes. Mulberry tree bark hydrolysate yielded the highest total sugars (16.4 g/L), but sessile oak bark hydrolysate showed the highest β -carotene production (362.7 mg/L) after 30 hours of fermentation. Woody biomasses are known to contain significant amounts of extractive inhibitory compounds. Surprisingly, when we removed them in order to promote growth and production, we observed a significant drop in carotenoids titers, which resulted in line with published productions from biomasses lacking extractive components. These data suggest that stressful compounds present in the extractive fractions are crucial for promoting high production and productivity, when compared with the use of biomasses lacking such components. This research therefore highlights the potential of underexplored urban woody residues, thanks to the presence of triggering components, as an advantageous feedstock for microbial carotenoids production.